Author Index

Abbott, F., see Kassahun, K. 327
Abernethy, D.R., see DePetrillo, P.B. 421
Abramson, F.P., see Chace, D.H. 1
Airoldi, L., see Bagnati, R. 267
Ambroise, C., see DePetrillo, P.B. 421
Andrew, R., see Midgley, J.M. 259
Arrigoni Martelli, E., see Marzo, A. 247
Atkinson, S.A., see McPhee, M.D. 41
Austin, D.
—, Koziol, T., Woolf, E. and Espinosa-Crouch, G.
Normal-phase high-performance liquid chromatographic procedure for the determination of diprafenone in plasma 182
Bagnati, R.
—, Grazia Castelli, M., Airoldi, L., Paleologo Oriundi, M., Ubaldi, A. and Fanelli, R.
Analysis of diethylstilbestrol, dienestrol and hexestrol in biological samples by immunoaffinity extraction and gas chromatography–negative-ion chemical ionization mass spectrometry 267
Basci, N.E.
—, Bozkurt, A., Kayaalp, S.O. and Isimer, A.
Comparison of colorimetric and high-performance liquid chromatographic determination of sulphamethoxazole and acetylsuphamethoxazole 174
Behrens, R., see Taylor, R.B. 490
Beyne, P.
—, Doute, M., Lacour, B., Lederer, F. and Pays, M.
Simple and rapid method for the purification of human apolipoprotein C II by reversed-phase high-performance liquid chromatography 140
Bombardt, P.A.
—, Brewer, J.E., Thomas, Jr., R.C. and Koets, M.D.
Simple, rapid and sensitive high-performance liquid chromatographic procedure for the quantitation of itazigrel in human plasma or serum 233
Bonati, M., see Celardo, A. 220
Bozkurt, A., see Basci, N.E. 174
Brewer, J.E., see Bombardt, P.A. 233
Bus, C.A., see Norwood, D.L. 289
Cardace, G., see Marzo, A. 247
Carter, A.J.
— and Müller, R.E.
Application and validation of an ion-exchange high-performance liquid chromatographic method for measuring adenine nucleotides, creatine and creatine phosphate in mouse brain 31
Castelli, M. Grazia, see Bagnati, R. 267
Celardo, A.
— and Bonati, M.
Determination of thiopental measured in human blood by reversed-phase high-performance liquid chromatography 220
Chace, D.H.
— and Abramson, F.P.
Detection of carbon-14 eluting from a capillary gas chromatograph using chemical reaction interface–mass spectrometry 1
Chaumont, A.J., see Kennel, S. 169
Cheng, L.K., see Greene, S.V. 397
Churchill, F.C., see Mount, D.L. 51
—, see Wages, S.A. 115
Clifford, A.J., see Furr, H.C. 406
Cole, D.E.C., see McPhee, M.D. 41
Corasaniti, M.T.
—, Strongoli, M.C. and Nisticò, G.
Determination of paraquat in rat brain using ion-pair solid-phase extraction and reversed-phase high-performance liquid chromatography with ultraviolet detection 189
Dandekar, K.A., see Mulvana, D.E. 343
Dawson, C.M., see Rainbow, S.J. 389
Decourt, J.P.
—, Mura, P., Papet, Y., Piriou, A. and Reiss, D.
Simultaneous determination of progabide and its acid metabolite by reversed-phase high-performance liquid chromatography 214
Dell, D., see Heizmann, P. 91
DePetrillo, P.B.
—, Swift, R.M., Ambroise, C. and Abernethy, D.R.
High-performance liquid chromatographic determination of 3′,5′-cyclic adenosine monophosphate in human platelets 421

Derendorf, H., see Schaefer, H.G. 351
Doute, M., see Beyne, P. 140
Doyle, E.
—, McDowall, R.D., Murkitt, G.S., Picot, V.S. and Rogers, S.J.
Two systems for the automated analysis of drugs in biological fluids using high-performance liquid chromatography 67
—, see Picot, V.S. 454
Doyle, M.J., see Kaffenberger, R.M. 59
Durand, A., see Rop, P.P. 226
Eggers, H., see Heizmann, P. 91
Eigendorf, G.K., see Salari, H. 303
Engineer, M.S.
— and Newman, R.A.
High-performance liquid chromatographic measurement of (*R*)-4-[3-[(2-hydroxy-2-phenethyl)amino]-3-methylbutyl]benzamide monohydrochloride (LY195448) and its *p*-hydroxy metabolite in human plasma and urine 473
Eriksson, B.-M.
—, Persson, B.-A. and Wikström, M.
Determination of urinary vanillylmandelic acid by direct injection and coupled-column chromatography with electrochemical detection 11
Espinosa-Crouch, G., see Austin, D. 182
Fanelli, R., see Bagnati, R. 267
Farrell, K., see Kassahun, K. 327
Ferraris, S.
— and Korte, Jr., D.W.
Determination of the oximes 2-hydroxyiminomethyl-3-methyl-1-[2-(3-methyl-3-nitrobutyloxymethyl)]imidazolium chloride and 1-[1-(3-butynyloxymethyl)]-2-hydroxyiminomethyl-3-methylimidazolium chloride in plasma by high-performance liquid chromatography 361
Foth, H., see Michaelis, H.C. 201
Fukasawa, K., see Harada, M. 158
Fukasawa, K.M., see Harada, M. 158
Furr, H.C.
—, Zeng, S., Clifford, A.J. and Olson, J.A.
Capillary gas chromatography of retinoids (vitamin A compounds) and apo-retinoids: determination of Kováts retention indices 406
Geng, W., see Michaelis, H.C. 201
Glennon, J.D.
— and Senior, A.T.
Simultaneous determination of desferrioxamine and ferrioxamine by high-performance liquid chromatography with amperometric detection 481
Gora, R., see Heizmann, P. 91
Grazia Castelli, M., see Bagnati, R. 267
Greene, S.V.
—, Krebs, H.A., Lemieux, R.E. and Cheng, L.K.
Determination of a new anti-allergenic agent, 1-[4-[3-[4-[bis-(4-fluorophenyl)hydroxymethyl]-1-piperidinyl]-propoxy]-3-methoxyphenyl]ethanone, and its active acidic metabolite in plasma by high-performance liquid chromatography 397
Gristwood, W.E.
Determination of fluparoxan (GR50360) in plasma by gas chromatography 436
Grossklaus, R., see Klingebiel, L. 238
Ha, P.T., see Volmut, J. 428
Harada, M.
—, Mogi, M., Hiraoka, B.Y., Fukasawa, K., Fukasawa, K.M., Takagi, T. and Sasaki, S.
Determination of proline iminopeptidase activity in bovine dental germs by high-performance liquid chromatography 158
Hashimoto, Y., see Nakanuma, R. 208
Heizmann, P.
—, Dell, D., Eggers, H. and Gora, R.
Determination of the new fluoroquinolone fleroxacin and its N-demethyl and N-oxide metabolites in plasma and urine by high-performance liquid chromatography with fluorescence detection 91
Henke, W.
—, Nickel, E. and Jung, K.
Determination of purine compounds by ion-pair microbore high-performance liquid chromatography: application to ischemic rat kidney mitochrondria 498
Heyworth, P., see Leoni, P. 152
Hiraoka, B.Y., see Harada, M. 158
Ho, J.W.
Micro assay for urinary δ-aminolevulinic acid and porphobilinogen by high-performance liquid chromatography with pre-column derivatization 134
Horiuchi, M., see Morishima, K. 381

Houen, G.
Affinity chromatographic isolation of a major polyamine binding protein from serum and placenta, identified as serum albumin 146
Ikemoto, S., see Isoda, N. 315
Isimer, A., see Basci, N.E. 174
Isoda, N.
—, Kajii, E., Ikemoto, S. and Kimura, K.
Esophageal carcinoma-associated proteins detected by two-dimensional polyacrylamide gel electrophoresis 315
Jørgensen, A., see Rop, P.P. 226
Jung, K., see Henke, W. 498
Kadir, F.
— and Zuidema, J.
High-performance liquid chromatographic method for the determination of carazolol in serum or plasma of pigs 461
Kaffenberger, R.M.
— and Doyle, M.J.
Determination of menthol and menthol glucuronide in human urine by gas chromatography using an enzyme-sensitive internal standard and flame ionization detection 59
Kågedal, B.
— and Olsson, B.
Determination of creatinine in serum by high-performance liquid chromatography: a comparison of three ion-exchange methods 21
Kahl, G.F., see Michaelis, H.C. 201
Kajii, E., see Isoda, N. 315
Kassahun, K.
—, Farrell, K., Zheng, J. and Abbott, F.
Metabolic profiling of valproic acid in patients using negative-ion chemical ionization gas chromatography–mass spectrometry 327
Kaul, S., see Mulvana, D.E. 343
Kayaalp, S.O., see Basci, N.E. 174
Kennel, S.
—, Kintz, P., Tracqui, A., Mangin, P., Lugnier, A.A. and Chaumont, A.J.
Identification and quantification in plasma of zopiclone by gas chromatography with nitrogen–phosphorus detection 169
Kimura, K., see Isoda, N. 315
Kintz, P., see Kennel, S. 169
Klingebiel, L.
—, Krüger, D. and Grosasklaus, R.
Analysis of Palatinit® (isomalt) and its monomers in rat intestinal samples by high-performance liquid chromatography with ultraviolet detection 238
Kodama, H.
—, Nakamura, H., Sugahara, K. and Numajiri, Y.
Liquid chromatography–mass spectrometry for the qualitative analyses of iminodipeptides in the urine of patients with prolidase deficiency 279
Koets, M.D., see Bombardt, P.A. 233
Korte, Jr., D.W., see Ferraris, S. 361
Koziol, T., see Austin, D. 182
Kramer, H.K., see Wiener, H.L. 467
Krebs, H.A., see Greene, S.V. 397
Krüger, D., see Klingebiel, L. 238
Lachno, D.R., see Smolenski, R.T. 414
Lacour, B., see Beyne, P. 140
Lederer, F., see Beyne, P. 140
Ledingham, S.J.M., see Smolenski, R.T. 414
Lemieux, R.E., see Greene, S.V. 397
Leoni, P.
—, Heyworth, P. and Segal, A.W.
Separation of phosphoproteins by fast protein liquid chromatography 152
Lopez, L.M., see Schaefer, H.G. 351
Lugnier, A.A., see Kennel, S. 169
MacDonald, N., see Midgley, J.M. 259
McDowall, R.D., see Doyle, E. 67
McPhee, M.D.
—, Atkinson, S.A. and Cole, D.E.C.
Quantitation of free sulfate and total sulfoesters in human breast milk by ion chromatography 41
Mangin, P., see Kennel, S. 169
Martelli, E. Arrigoni, see Marzo, A. 247
Marzo, A.
—, Cardace, G., Monti, N., Muck, S. and Arrigoni Martelli, E.
Chromatographic and non-chromatographic assay of L-carnitine family components 247
Matisová, E., see Volmut, J. 428
Mehvar, R.
Liquid chromatographic analysis of propafenone enantiomers in human plasma 79
Michaelis, H.C.
—, Geng, W., Kahl, G.F. and Foth, H.
Sensitive determination of bupivacaine

in human plasma by high-performance liquid chromatography 201
Midgley, J.M.
—, Andrew, R., Watson, D.G., MacDonald, N., Reid, J.L. and Williams, D.A.
Dichlorotetrafluoroacetone as a derivatisation reagent in the analysis of mandelic acids in human urine 259
Millington, D.S., see Norwood, D.L. 289
Mizuguchi, S., see Nakanuma, R. 208
Mogi, M., see Harada, M. 158
Monti, N., see Marzo, A. 247
Moody, R.R., see Taylor, R.B. 490
Morishima, K.
—, Tahara, R. and Horiuchi, M.
Determination of a new calcium antagonist, sesamodil fumarate (SD-3211), and its metabolite in plasma by liquid chromatography with electrochemical detection 381
Morita, I.
— and Yoshida, H.
Direct injection of highly protein-bound compounds by column-switching high-performance liquid chromatography 127
Motohashi, M., see Yamashita, K. 103, 196
Mount, D.L.
—, Patchen, L.C. and Churchill, F.C.
Determination of mefloquine in blood by supercritical fluid chromatography with electron-capture detection 51
Muck, S., see Marzo, A. 247
Müller, R.E., see Carter, A.J. 31
Mulvana, D.E.
—, Kaul, S., Dandekar, K.A. and Pittman, K.A.
Determination of belfosdil, a new calcium channel blocker, in human plasma by capillary gas chromatography with nitrogen–phosphorus detection 343
Mura, P., see Decourt, J.P. 214
Murkitt, G.S., see Doyle, E. 67
Nakamura, H., see Kodama, H. 279
Nakamura, T., see Nakanuma, R. 208
Nakanuma, R.
—, Watanabe, K., Yamashita, K., Mizuguchi, S., Hashimoto, Y., Nakamura, T. and Umezawa, H.
High-performance liquid chromatographic determination of deoxyspergualin in dog plasma with ultraviolet detection 208
Nambara, T., see Ohkubo, T. 441
Newman, R.A., see Engineer, M.S. 473
Nickel, E., see Henke, W. 498
Nisticò, G., see Corasaniti, M.T. 189
Norwood, D.L.
—, Bus, C.A. and Millington, D.S.
Combined high-performance liquid chromatographic–continuous-flow fast atom bombardment mass spectrometric analysis of acylcoenzyme A compounds 289
Numajiri, Y., see Kodama, H. 279
Ohkubo, T.
— and Nambara, T.
Determination of mitomycin C in serum by high-performance liquid chromatography with dual-electrode coulometric detection 441
Olson, J.A., see Furr, H.C. 406
Olsson, B., see Kågedal, B. 21
Oriundi, M. Paleologo, see Bagnati, R. 267
Page, R.L., see Tyczkowska, K. 447
Paleologo Oriundi, M., see Bagnati, R. 267
Papet, Y., see Decourt, J.P. 214
Patchen, L.C., see Mount, D.L. 51
—, see Wages, S.A. 115
Paya, M., see Beyne, P. 140
Pearce, J.C., see Picot, V.S. 454
Persson, B.A., see Eriksson, B.-M. 11
Picot, V.S.
—, Doyle, E. and Pearce, J.C.
Analysis of Zaprinast in rat and human plasma by automated solid-phase extraction and reversed-phase high-performance liquid chromatography 454
—, see Doyle, E. 67
Piriou, A., see Decourt, J.P. 214
Pittman, K.A., see Mulvana, D.E. 343
Rainbow, S.J.
—, Dawson, C.M. and Tickner, T.R.
Direct serum injection high-performance liquid chromatographic method for the simultaneous determination of phenobarbital, carbamazepine and phenytoin 389
Reid, J.L., see Midgley, J.M. 259
Reiss, D., see Decourt, J.P. 214
Reith, M.E.A., see Wiener, H.L. 467
Riviere, J.E., see Tyczkowska, K. 447
Rogers, S.J., see Doyle, E. 67

Rop, P.P.
—, Durand, A., Viala, A. and Jørgensen, A.
Simultaneous determination of citalopram, monodesmethylcitalopram and didesmethylcitalopram in plasma by high-performance liquid chromatography after column extraction 226
Salari, H.
— and Eigendorf, G.K.
Detection of lyso-platelet-activating factor by high-performance liquid chromatography after derivatisation with fluorescent fatty acids 303
Salem, G.W., see Shih, M.L. 371
Salonen, J.S.
Determination of the amine metabolites of selegiline in biological fluids by capillary gas chromatography 163
Sasaki, S., see Harada, M. 158
Schaefer, H.G.
—, Spahn, H., Lopez, L.M. and Derendorf, H.
Simultaneous determination of propranolol and 4-hydroxypropranolol enantiomers after chiral derivatization using reversed-phase high-performance liquid chromatography 351
Segal, A.W., see Leoni, P. 152
Senior, A.T., see Glennon, J.D. 481
Shih, M.L.
—, Smith, J.R. and Salem, G.W.
High-performance liquid chromatographic analysis of imidazolium and pyridinium oximes in plasma and urine 371
Smith, J.R., see Shih, M.L. 371
Smolenski, R.T.
—, Lachno, D.R., Ledingham, S.J.M. and Yacoub, M.H.
Determination of sixteen nucleotides, nucleosides and bases using high-performance liquid chromatography and its application to the study of purine metabolism in hearts for transplantation 414
Spahn, H., see Schaefer, H.G. 351
Statler, J.A.
Determination of tobramycin using high-performance liquid chromatography with pulsed amperometric detection 244
Strongoli, M.C., see Corasaniti, M.T. 189
Sugahara, K., see Kodama, H. 279
Swift, R.M., see DePetrillo, P.B. 421
Tahara, R., see Morishima, K. 381
Takagi, T., see Harada, M. 158
Taylor, R.B.
—, Behrens, R., Moody, R.R. and Wangboonskul, J.
Assay method for the simultaneous determination of proguanil, chloroquine and their major metabolites in biological fluids 490
Thomas, Jr., R.C., see Bombardt, P.A. 233
Ticknir, T.R., see Rainbow, S.J. 389
Tracqui, A., see Kennel, S. 169
Tyczkowska, K.
—, Page, R.L. and Riviere, J.E.
Determination of carboplatin in canine plasma by liquid chromatography with ultraviolet–visible detection and confirmation by atomic absorption spectroscopy 447
Ubaldi, A., see Bagnati, R. 267
Umezawa, H., see Nakanuma, R. 208
Viala, A., see Rop, P.P. 226
Volmut, J.
—, Matisová, E. and Ha, P.T.
Simultaneous determination of six antiepileptic drugs by capillary gas chromatography 428
Wages, S.A.
—, Patchen, L.C. and Churchill, F.C.
Analysis of blood and urine samples from *Macaca mulata* for pyronaridine by high-performance liquid chromatography with electrochemical detection 115
Wangboonskul, J., see Taylor, R.B. 490
Watanabe, K., see Nakanuma, R. 208
Watson, D.G., see Midgley, J.M. 259
Wiener, H.L.
—, Kramer, H.K. and Reith, M.E.A.
Separation and determination of sertraline and its metabolite, desmethylsertraline, in mouse cerebral cortex by reversed-phase high-performance liquid chromatography 467
Wikström, M., see Eriksson, B.-M. 11
Williams, D.A., see Midgley, J.M. 259
Woolf, E., see Austin, D. 182
Yacoub, M.H., see Smolenski, R.T. 414
Yamashita, K.
—, Motohashi, M. and Yashiki, T.
High-performance liquid chromatographic determination of phenylpropanolamine in human plasma and urine, us-

ing column switching combined with ion-pair chromatography 103
—, Motohashi, M. and Yashiki, T. Sensitive high-performance liquid chromatographic determination of propranolol in human plasma with ultraviolet detection using column switching combined with ion-pair chromatography 196
—, see Nakanuma, R. 208
Yashiki, T., see Yamashita, K. 103, 196
Yoshida, H., see Morita, I. 127
Zeng, S., see Furr, H.C. 406
Zheng, J., see Kassahun, K. 327
Zuidema, J., see Kadir, F. 461

Subject Index

Acetyl-L-carnitine
Chromatographic and non-chromatographic assay of L-carnitine family components 247
Acetylsulphamethoxazole
Comparison of colorimetric and HPLC determination of sulphamethoxazole and acetylsulphamethoxazole 174
Acid progabide
Simultaneous determination of progabide and its acid metabolite by reserved-phase HPLC 214
Acylcoenzyme A
Combined HPLC–continuous-flow FAB MS analysis of acylcoenzyme A compounds 289
Adenine nucleotides
Application and validation of an anion-exchange HPLC method for measuring adenine nucleotides, creatine and creatine phosphate in mouse brain 31
Adenosine phosphates
Application and validation of an anion-exchange HPLC method for measuring adenine nucleotides, creatine and creatine phosphate in mouse brain 31
Albumin
Affinity chromatographic isolation of a major polyamine binding protein from serum and placenta, identified as serum albumin 146
Aminoglycosides
Determination of tobramycin using HPLC with pulsed amperometric detection 244
δ-Aminolevulinic acid
Micro assay for urinary δ-aminolevulinic acid and porphobilinogen by HPLC with pre-column derivatization 134
Amphetamine
Determination of the amine metabolites of selegiline in biological fluids by capillary GC 163
Antibiotics
Determination of tobramycin using HPLC with pulsed amperometric detection 244
Antiepileptic drugs
Simultaneous determination of six antiepileptic drugs by capillary GC 428
Apolipoprotein C II
Simple and rapid method for the purification of human apolipoprotein C II by reversed-phase HPLC 140
Apo-retinoids
Capillary GC of retinoids (vitamin A compounds) and apo-retinoids: determination of Kováts retention indices 406
Belfosdil
Determination of belfosdil, a new calcium channel blocker, in human plasma by capillary GC with nitrogen-phosphorus detection 343
1-[4-[3-[4-[Bis-(4-fluorophenyl)hydroxymethyl]-1-piperidinyl]propoxy]-3-methoxyphenyl]ethanone
Determination of a new anti-allergenic agent, 1-[4-[3-[4-[bis-(4-fluorophenyl) hydroxymethyl]-1-piperidinyl]propoxy] 3-methoxyphenyl]ethanone, and its active acidic metabolite in plasma by HPLC 397
Bupivacaine
Sensitive determination of bupivacaine in human plasma by HPLC 201
1-[1-(3-Butynyloxymethyl)]-2-hydroxyiminomethyl-3-methylimidazolium chloride
Determination of the oximes 2-hydroxyiminomethyl-3-methyl-1-[2-(3-methyl-3-nitrobutyloxymethyl)]imidazolium chloride and 1-[1-(3-butynyloxymethyl)]-2-hydroxyiminomethyl-3-methylimidazolium chloride in plasma by HPLC 361
Carazolol
HPLC method for the determination of carazolol in serum or plasma of pigs 461
Carbamazepine
Direct serum injection HPLC method for the simultaneous determination of phenobarbital, carbamazepine and phenytoin 389
Carbamazepine
Simultaneous determination of six antiepileptic drugs by capillary GC 428
Carbon-14
Detection of carbon-14 eluting from a cap-

illary gas chromatograph using chemical reaction interface–MS 1

Carboplatin
Determination of carboplatin in canine plasma by LC with UV–visible detection and confirmation by atomic absorption spectroscopy 447

L-Carnitine
Review. Chromatographic and non-chromatographic assay of L-carnitine family components 247

Chloroquine
Assay method for the simultaneous determination of proguanil, chloroquine and their major metabolites in biological fluids 490

Citalopram
Simultaneous determination of citalopram, monodesmethylcitalopram and didesmethylcitalopram in plasma by HPLC after column extraction 226

Coenzyme A thioesters
Combined HPLC–continuous-flow FAB MS analysis of acylcoenzyme A compounds 289

Creatine
Application and validation of an anion-exchange HPLC method for measuring adenine nucleotides, creatine and creatine phosphate in mouse brain 31

Creatine phosphate
Application and validation of an anion-exchange HPLC method for measuring adenine nucleotides, creatine and creatine phosphate in mouse brain 31

Creatinine
Determination of creatinine in serum by HPLC: a comparison of three ion-exchange methods 21

3′,5′-Cyclic adenosine monophosphate
HPLC determination of 3′,5′-cyclic adenosine monophosphate in human platelets 421

N-Demethylfleroxacin
Determination of the new fluoroquinolone fleroxacin and its N-demethyl and N-oxide metabolites in plasma and urine by HPLC with fluorescence detection 91

Deoxyspergualin
HPLC determination of deoxyspergualin in dog plasma with UV detection 208

Desferrioxamine
Simultaneous determination of desferrioxamine and ferrioxamine by HPLC with amperometric detection 481

Desmethylselegiline
Determination of the amine metabolites of selegiline in biological fluids by capillary GC 163

Desmethylsertraline
Separation and determination sertraline and its metabolite, desmethylsertraline, in mouse cerebral cortex by reversed-phase HPLC 467

Dichlorotetrafluoroacetone
Dichlorotetrafluoroacetone as a derivatisation reagent in the analysis of mandelic acid in human urine 259

Didesmethylcitalopram
Simultaneous determination of citalopram, monodesmethylcitalopram and didesmethylcitalopram in plasma by HPLC after column extraction 226

Dienestrol
Analysis of diethylstilbestrol, dienestrol and hexestrol in biological samples by immunoaffinity extraction and GC–negative-ion chemical ionization MS 267

Diethylstilbestrol
Analysis of diethylstilbestrol, dienestrol and hexestrol in biological samples by immunoaffinity extraction and GC–negative-ion chemical ionization MS 267

1,1′-Dimethyl-4,4′-bipyridinium dichloride
Determination of paraquat in rat brain using ion-pair solid-phase extraction and reversed-phase HPLC with UV detection 189

Diprafenone
Normal-phase HPLC procedure for the determination of diprafenone in plasma 182

Enantiomer separation
LC analysis of propafenone enantiomers in human plasma 79

Enzymes
Determination of proline iminopeptidase activity in bovine dental germs by HPLC 158

Esophagael carcinoma-associated proteins
Esophagael carcinoma-associated proteins detected by two-dimensional poly-

acrylamide gel electrophoresis 315
Estrogens
Analysis of diethylstilbestrol, dienestrol and hexestrol in biological samples by immunoaffinity extraction and GC–negative-ion chemical ionization MS 267
Ethosuximide
Simultaneous determination of six antiepileptic drugs by capillary GC 428
Ferrioxamine
Simultaneous determination of desferrioxamine and ferrioxamine by HPLC with amperometric detection 481
Fleroxacin
Determination of the new fluoroquinolone fleroxacin and its N-demethyl and N-oxide metabolites in plasma and urine by HPLC with fluorescence detection 91
Fluoroquinolone
Determination of the new fluoroquinolone fleroxacin and its N-demethyl and N-oxide metabolites in plasma and urine by HPLC with fluorescence detection 91
Fluoroquinolone N-oxide
Determination of the new fluoroquinolone fleroxacin and its N-demethyl and N-oxide metabolites in plasma and urine by HPLC with fluorescence detection 91
Fluparoxan
Determination of fluparoxan (GR50360) in plasma by GC 436
Glucose
Analysis of Palatinit® (isomalt) and its monomers in rat intestinal samples by HPLC with UV detection 238
GR50360
Determination of fluparoxan (GR50360) in plasma by GC 436
Hexestrol
Analysis of diethylstilbestrol, dienestrol and hexestrol in biological samples by immunoaffinity extraction and GC–negative-ion chemical ionization MS 267
H^+/K^+ ATPase inhibitor
Two systems for the automated analysis of drugs in biological fluids using HPLC 67
2-Hydroxyiminomethyl-3-methyl-1-[2-(3-methyl-3-nitrobutyloxymethyl)]imidazolium chloride
Determination of the oximes 2-hydroxyiminomethyl-3-methyl-1-[2-(3-methyl-3-nitrobutyloxymethyl)]imidazolium chloride and 1 1-[1-(3-butynyloxymethyl)]-2-hydroxyiminomethyl-3-methylimidazolium chloride in plasma by HPLC 361
Hydroxymandelic acids
Dichlorotetrafluoroacetone as a derivatisation reagent in the analysis of mandelic acids in human urine 259
(*R*)-4-[3-[(2-Hydroxy-2-phenethyl)amino]-3-methylbutyl]benzamide monohydrochloride
HPLC measurement of (*R*)-4-[3-[(2-hydroxy-2-phenethyl)amino]-3-methylbutyl]benzamide monohydrochloride (LY195448) and its *p*-hydroxy metabolite in human plasma and urine 473
4-Hydroxypropranolol enantiomers
Simultaneous determination of propranolol and 4-hydroxypropranolol enantiomers after chiral derivatization using reversed phase HPLC 351
Imidazolium oximes
HPLC analysis of imidazolium and pyridinium oximes in plasma and urine 371
Iminodipeptides
LC–MS for the qualitative analyses of iminodipeptides in the urine of patients with prolidase deficiency 279
Indoles
Direct injection of highly protein-bound compounds by column-switching HPLC 127
Isomalt
Analysis of Palatinit® (isomalt) and its monomers in rat intestinal samples by HPLC with UV detection 238
Itazigrel
Simple, rapid and sensitive HPLC procedure for the quantitation of itazigrel in human plasma or serum 233
LY195448
HPLC measurement of (*R*)-4-[3-[(2-hydroxy-2-phenethyl)amino]-3-methylbutyl]benzamide monohydrochloride (LY195448) and its *p*-hydroxy metabolite in human plasma and urine 473
Lyso-platelet-activating factor
Detection of lyso-platelet-activating factor by HPLC after derivatisation with fluorescent fatty acids 303

Mandelic acids
Dichlorotetrafluoroacetone as a derivatisation reagent in the analysis of mandelic acids in human urine 259
Mannitol
Analysis of Palatinit® (isomalt) and its monomers in rat intestinal samples by HPLC with UV detection 238
Mefloquine
Determination of mefloquine in blood by supercritical fluid chromatography with electron-capture detection 51
Menthol
Determination of menthol and menthol glucuronide in human urine by GC using an enzyme-sensitive internal standard and flame ionization detection 59
Menthol glucuronide
Determination of menthol and menthol glucuronide in human urine by GC using an enzyme-sensitive internal standard and flame ionization detection 59
Metamphetamine
Determination of the amine metabolites of selegiline in biological fluids by capillary GC 163
N-Methyloleoyl taurate
Determination of creatinine in serum by HPLC: a comparison of three ion-exchange methods 21
Mitomycin C
Determination of mitomycin C in serum by HPLC with dual-electrode coulometric detection 441
Monodesmethylcitalopram
Simultaneous determination of citalopram, monodesmethylcitalopram and didesmethylcitalopram in plasma by HPLC after column extraction 226
Nucleobases
Determination of sixteen nucleotides, nucleosides and bases using HPLC and its application to the study of purine metabolism in hearts for transplantation 414
Nucleobases
Determination of purine compounds by ion-pair microbore HPLC: application to ischemic rat kidney mitochondria 498
Nucleosides
Determination of sixteen nucleotides, nucleosides and bases using HPLC and its application to the study of purine metabolism in hearts for transplantation 414
Nucleosides
Determination of purine compounds by ion-pair microbore HPLC: application to ischemic rat kidney mitochondria 498
Nucleotides
Application and validation of an anion-exchange HPLC method for measuring adenine nucleotides, creatine and creatine phosphate in mouse brain 31
Nucleotides
Determination of sixteen nucleotides, nucleosides and bases using HPLC and its application to the study of purine metabolism in hearts for transplantation 414
Nucleotides
Determination of purine compounds by ion-pair microbare HPLC: application to ischemic rat kidney mitochondria 498
Oximes
Determination of the oximes 2-hydroxyiminomethyl-3-methyl-1-[2-(3-methyl-3-nitrobutyloxymethyl)]imidazolium chloride and 1-[1-(3-butynyloxymethyl)]-2-hydroxyiminomethyl-3-methylimidazolium chloride in plasma by HPLC 361
Oximes
HPLC analysis of imidazolium and pyridinium oxides in plasma and urine 371
Palatinit®
Analysis of Palatinit® (isomalt) and its monomers in rat intestinal samples by HPLC with UV detection 238
Paraquat
Determination of paraquat in rat brain using ion-pair solid-phase extraction and reversed-phase HPLC with UV detection 189
Peptides
LC–MS for the qualitative analyses of iminodipeptides in the urine of patients with prolidase deficiency 279
Phenobarbital
Direct serum injection HPLC method for the simultaneous determination of phenobarbital, carbamazepine and phenytoin 389
Phenobarbital
Simultaneous determination of six antiepileptic drugs by capillary GC 428

Phenylpropanolamine
HPLC determination of phenylpropanolamine in human plasma and urine, using column switching combined with ion-pair chromatography 103
Phenytoin
Detection of carbon-14 eluting from a capillary gas chromatograph using chemical reaction interface–MS 1
Phenytoin
Direct serum injection HPLC method for the simultaneous determination of phenobarbital, carbamazepine and phenytoin 389
Phenytoin
Simultaneous determination of six antiepileptic drugs by capillary GC 428
Phosphoproteins
Separation of phosphoproteins by fast protein LC 152
Platelet-activating factor
Detection of lyso-platelet-activating factor by HPLC after derivatisation with fluorescent fatty acids 303
Polyamines
Affinity chromatographic isolation of a major polyamine binding protein from serum and placenta, identified as serum albumin 146
Porphobilinogen
Micro assay for urinary δ-aminolevulinic acid and porphobilinogen by HPLC with pre-column derivatization 134
Primidone
Simultaneous determination of six antiepileptic drugs by capillary GC 428
Progabide
Simultaneous determination of progabide and its acid metabolite by reversed-phase HPLC 214
Proguanil
Assay method for the simultaneous determination of proguanil, chloroquine and their major metabolites in biological fluids 490
Proline iminopeptidase
Determination of proline iminopeptidase activity in bovine dental germs by HPLC 158
Propafenone
LC analysis of propafenone enantiomers in human plasma 79
Propionyl-L-carnitine
Review. Chromatographic and non-chromatographic assay of L-carnitine family components 247
Propranolol
Sensitive HPLC determination of propranolol in human plasma with UV detection using column switching combined with ion-pair chromatography 196
Propranolol enantiomers
Simultaneous determination of propranolol and 4-hydroxypropranolol enantiomers after chiral derivatization using reversed-phase HPLC 351
Protein-bound compounds
Direct injection of highly protein-bound compounds by column-switching HPLC 127
Proteins
Affinity chromatographic isolation of a major polyamine binding protein from serum and placenta, identified as serum albumin 146
Proteins
Separation of phosphoproteins by fast protein LC 152
Proteins
Esophageal carcinoma-associated proteins detected by two-dimensional polyacrylamide gel electrophoresis 315
Purine compounds
Determination of purine compounds by ion-pair microbore HPLC: application to ischemic rat kidney mitochondria 498
Pyridinium oximes
HPLC analysis of imidazolium and pyridinium oximes in plasma and urine 371
Pyronaridine
Analysis of blood and urine samples from *Macaca mulata* for pyronaridine by HPLC with electrochemical detection 115
Retinoids
Capillary GC of retinoids (vitamin A compounds) and apo-retinoids: determination of Kováts retention indices 406
Sample preparation
Two systems for the automated analysis of drugs in biological fluids using HPLC 67

SD-3211
Determination of a new calcium antagonist, sesamodil fumarate (SD-3211), and its metabolite in plasma by LC with electrochemical detection 381
Selegiline
Determination of the amine metabolites of selegiline in biological fluids by capillary GC 163
Sertraline
Separation and determination of sertraline and its metabolite, desmethylsertraline, in mouse cerebral cortex by reversed-phase HPLC 467
Sesamodil fumarate
Determination of a new calcium antagonist, sesamodil fumarate (SD-3211), and its metabolite in plasma by LC with electrochemical detection 381
Sorbitol
Analysis of Palatinit® (Isomalt) and its monomers in rat intestinal samples by HPLC with UV detection 238
Steroids
Analysis of diethylstilbestrol, dienestrol and hexestrol in biological samples by immunoaffinity extraction and GC–negative-ion chemical ionization MS 267
Sulphamethoxazole
Comparison of colorimetric and HPLC determination of sulphamethoxazole and acetylsulphamethoxazole 174
Sulphate
Quantitation of free sulfate and total sulfoesters in human breast milk by ion chromatography 41
Sulpho esters
Quantitation of free sulfate and total sulfoesters in human breast milk by ion chromatography 41
Supercritical fluid chromatography
Determination of mefloquine in blood by supercritical fluid chromatography with electron-capture detection 51
Thiopental
Determination of thiopental measured in human blood by reversed-phase HPLC 220
Tobramycin
Determination of tobramycin using HPLC with pulsed amperometric detection 244
Valproic acid
Metabolic profiling of valproic acid in patients using negative-ion chemical ionization GC–MS 327
Valproic acid
Simultaneous determination of six antiepileptic drugs by capillary GC 428
Vanillylmandelic acid
Determination of urinary vanilylmandelic acid by direct injection and coupled-column chromatography with electrochemical detection 11
Vanillylmandelic acid
Dichrlorotetrafluoroacetone as a derivatisation reagent in the analysis of mandelic acids in human urine 259
Vitamin A compounds
Capillary GC of retinoids (vitamin A compounds) and apo-retinoids: determination of Kováts retention indices 406
Zaprinast
Two systems for the automated analysis of drugs in biological fluids using HPLC 67
Zaprinast
Analysis of Zaprinast in rat and human plasma by automated solid-phase extraction and reversed-phase HPLC 454
Zopiclone
Identification and quantification in plasma of zopiclone by GC with nitrogen–phosphorus detection 169